

Name: Key Class: _____ Date: _____

ID: A

Trig Midterm Review 2015

Multiple Choice

Identify the letter of the choice that best completes the statement or answers the question.

Chapter 5
Starts
 $360 + 0.43(60)$
 $0.8(60)$

- B 1. Change 360.43° to degrees, minutes, and seconds.
- a. $360^\circ 55' 56''$
 - b. $360^\circ 25' 48''$
 - c. $360^\circ 56' 80''$
 - d. $360^\circ 53' 55''$

- D 2. Write $87^\circ 26' 3''$ as a decimal to the nearest thousandth.
- a. 87.437°
 - b. 87.444°
 - c. 87.484°
 - d. 87.434°

$87 + \frac{26}{60} + \frac{3}{3600}$

- C 3. Give the angle measure represented by 120° rotations clockwise.
- a. 43199°
 - b. -43203°
 - c. -43200°
 - d. 43201°


$120(-360)$

- C 4. Find the least positive angle measurement that is coterminal with -230° .
- a. 140°
 - b. 135°
 - c. 130°
 - d. 132°

Find the values of the sine, cosine, and tangent for $\angle A$.

A

5.




$4^2 + 7^2 = c^2$
 $\sqrt{65} = c$

- a. $\sin A = \frac{4\sqrt{65}}{65}, \cos A = \frac{7\sqrt{65}}{65}, \tan A = \frac{4}{7}$
- b. $\sin A = \frac{\sqrt{65}}{7}, \cos A = \frac{\sqrt{65}}{4}, \tan A = \frac{4}{7}$
- c. $\sin A = \frac{\sqrt{65}}{4}, \cos A = \frac{\sqrt{65}}{7}, \tan A = \frac{7}{4}$
- d. $\sin A = \frac{7\sqrt{65}}{65}, \cos A = \frac{4\sqrt{65}}{65}, \tan A = \frac{7}{4}$

B

6. If $\tan \theta = \frac{3}{4}$, find $\sin \theta$.



a. $\sin \theta = \frac{1}{2}$

c. $\sin \theta = 2$

b. $\sin \theta = \frac{3}{5}$

d. $\sin \theta = \frac{8}{5}$

D


7. Find $\cos \theta$ if θ is an angle in standard position and the point with coordinates $(-12, 5)$ lies on the terminal side of the angle.

a. $\frac{5}{13}$

c. $-\frac{13}{12}$

b. $-\frac{5}{12}$

d. $-\frac{12}{13}$



D

8. Find the values of the six trigonometric functions of an angle in standard position if the point with coordinates $(6, 8)$ lies on its terminal side.

a. $\sin \alpha = \frac{5}{4}, \cos \alpha = \frac{5}{3}, \tan \alpha = \frac{3}{4}$

c. $\sin \alpha = \frac{3}{5}, \cos \alpha = \frac{4}{5}, \tan \alpha = \frac{4}{3}$

$\csc \alpha = \frac{4}{5}, \sec \alpha = \frac{3}{5}, \cot \alpha = \frac{4}{3}$


$\csc \alpha = \frac{5}{3}, \sec \alpha = \frac{5}{4}, \cot \alpha = \frac{3}{4}$

b. $\sin \alpha = \frac{4}{3}, \cos \alpha = \frac{3}{4}, \tan \alpha = \frac{4}{5}$

d. $\sin \alpha = \frac{4}{5}, \cos \alpha = \frac{3}{5}, \tan \alpha = \frac{4}{3}$


$\csc \alpha = \frac{3}{4}, \sec \alpha = \frac{4}{3}, \cot \alpha = \frac{5}{4}$

$\csc \alpha = \frac{5}{4}, \sec \alpha = \frac{5}{3}, \cot \alpha = \frac{3}{4}$



B

9. If $g = 26.4$ and $F = 35^\circ$, find h . Round to the nearest tenth.



$$\cos 35 = \frac{h}{26.4}$$

a. $h = 22.6$

c. $h = 24.6$

b. $h = 21.6$

d. $h = 20.6$

Solve the equation if $0^\circ \leq x \leq 360^\circ$.

D

10. $\cos x = -\frac{1}{2}$

where on unit circle is $x = -\frac{1}{2}$

- a. $135^\circ, 225^\circ$
- b. $210^\circ, 330^\circ$

- c. $150^\circ, 210^\circ$
- d. $120^\circ, 240^\circ$

C

11. Name four angles whose tangent equals 0.

- a. $45^\circ, 135^\circ, 405^\circ, 495^\circ$
- b. $90^\circ, 270^\circ, 450^\circ, 630^\circ$

- c. $0^\circ, 180^\circ, 360^\circ, 540^\circ$
- d. $90^\circ, 450^\circ, 810^\circ, 1170^\circ$

B


12. Evaluate $\sec \left(\sin^{-1} \frac{\sqrt{3}}{2} \right)$. Assume that all the angles are in Quadrant I.

a. $\sqrt{3}$

c. $\frac{2\sqrt{3}}{3}$


b. 2

d. $\frac{1}{2}$



C

13. If $t = 38.5$ and $s = 31.4$, find S . Round to the nearest tenth.



$\sin S = \frac{31.4}{38.5}$

- a. $S = 53.6^\circ$
- b. $S = 56.6^\circ$

- c. $S = 54.6^\circ$
- d. $S = 55.6^\circ$

C

14. In right triangle ABC , $A = 28^\circ$, $b = 7$, and $\angle C$ is the right angle. Solve the triangle.

- a. $B = 62^\circ, a = 3.3, c = 7.7$
- b. $B = 62^\circ, a = 7.9, c = 3.7$

- c. $B = 62^\circ, a = 3.7, c = 7.9$
- d. $B = 62^\circ, a = 6.2, c = 9.4$

A

15. In right triangle ABC , $b = 6$, $c = 13$, and $\angle C$ is the right angle. Solve the triangle.

- a. $A = 63^\circ, B = 27^\circ, a = 11.5$
- b. $A = 27^\circ, B = 63^\circ, a = 11.5$

- c. $A = 65^\circ, B = 25^\circ, a = 14.3$
- d. $A = 25^\circ, B = 63^\circ, a = 14.3$

D

16. In right triangle ABC , $B = 75^\circ$, $c = 14$, and $\angle C$ is the right angle. Solve the triangle.

- a. $A = 15^\circ, a = 13.2, b = 4.7$
- b. $A = 15^\circ, a = 4.7, b = 13.2$


- c. $A = 15^\circ, a = 13.5, b = 3.7$
- d. $A = 15^\circ, a = 3.7, b = 13.5$

A


17. Given a triangle with $a = 19$, $A = 43^\circ$, and $B = 26^\circ$, what is the length of c ? Round to the nearest tenth.

- a. 26
- b. 25



- c. 27
- d. 28



$\frac{\sin 43}{19} = \frac{\sin 111}{c}$




$\tan 28 = \frac{a}{7}$



$\sin 15 = \frac{a}{19}$

Name: _____

$\frac{1}{2}(11)(8) \sin 60$



ID: A

B

18. Find the area of the triangle with $A = 60^\circ$, $b = 11$ feet, and $c = 8$ feet. Round to the nearest tenth.

- a. 22 ft^2
- b. 38.1 ft^2
- c. 44 ft^2
- d. 12.4 ft^2

D

19. How many triangles are there that satisfy the conditions $a = 14$, $b = 2$, $\alpha = 66^\circ$?


- a. impossible to determine
- b. 2
- c. 0
- d. 1

$A < 90^\circ \quad a > b \rightarrow 1 \Delta$

A

20. Given a triangle with $b = 7$, $c = 3$, and $A = 37^\circ$ what is the length of a ? Round to the nearest tenth.

- a. 4.9
- b. 5.9
- c. 5.5
- d. 4.3



B

21. Find the area of the triangle with $a = 18.8$, $b = 11.6$, $c = 13.8$. Round to the nearest tenth.

- a. 79.1 units^2
- b. 79.7 units^2
- c. 82.7 units^2
- d. 80.1 units^2

B

22. Change 1.96 radians to degree measure. Round to the nearest tenth.

- a. 472.3°
- b. 112.3°
- c. 292.3°
- d. 202.3°

$1.96 \left(\frac{180}{\pi} \right)$

B

23. Change 290° to radian measure in terms of π .

- a. $\frac{29}{27}\pi$
- b. $\frac{29}{18}\pi$
- c. $\frac{29}{36}\pi$
- d. $\frac{29}{9}\pi$

$\frac{290\pi}{180}$

C

24. Find the area of a sector with a central angle of 32° and a radius of 8.5 millimeters. Round to the nearest tenth.

- a. 40.4 mm^2
- b. 2.4 mm^2
- c. 20.2 mm^2
- d. 9.5 mm^2

$32 \cdot \frac{\pi}{180}$

D

25. A pulley of radius 10 cm turns at 6 revolutions per second. What is the linear velocity of the belt driving the pulley in meters per second?

- a. 376.99 m/s
- b. 1.67 m/s
- c. 166.67 m/s
- d. 3.77 m/s

$6(2\pi)$

A

26. Use a graph of the sine function to find the value of θ for which $\sin \theta = 0$.

- a. $\theta = \pi k$
- b. $\theta = \frac{\pi}{2} + 2\pi k$
- c. $\theta = \frac{\pi}{2} + \pi k$
- d. $\theta = 2\pi k$

$0.1(12\pi)$

Chapter 4 starts

B27. Find the amplitude, period, and phase shift of $f(x) = -4 \sin(7x + 2)$.

a. amplitude = -4

period = $\frac{2\pi}{7}$

phase shift = $\frac{2}{7}$

b. amplitude = 4

period = $\frac{2\pi}{7}$

phase shift = $-\frac{2}{7}$

c. amplitude = 8

period = $\frac{\pi}{7}$

phase shift = $-\frac{2}{7}$

d. amplitude = -4

period = 2π

phase shift = $\frac{2}{7}$

A28. Write an equation of the cosine function with amplitude 2 and period 4π .

a. $y = 2 \cos\left(\frac{1}{2}x\right)$

c. $y = -\frac{1}{2} \cos\left(\frac{1}{2}x\right)$

b. $y = -2 \cos\left(\frac{1}{4}x\right)$

d. $y = \frac{1}{2} \cos\left(\frac{1}{4}x\right)$

$$\frac{4\pi}{1} = \frac{2\pi}{K}$$


$$4\pi K = 2\pi$$

$$K = \frac{1}{2}$$

B


29. Graph the function. Which choice gives the amplitude, period, phase shift, and vertical shift for the function?

$$y = 4 \cos \left(3\theta + \frac{3}{4}\pi \right) + 4$$




4; $\frac{2}{3}\pi$; $-\frac{1}{4}\pi$; 4

b.




4; $\frac{2}{3}\pi$; $-\frac{1}{4}\pi$; 4



~~4; $\frac{2}{3}\pi$; $-\frac{1}{4}\pi$; 4~~

d.



~~4; $\frac{2}{3}\pi$; $-\frac{1}{4}\pi$; -4~~

D

30. Write an equation of the cosine function with the given amplitude, period, phase shift, and vertical shift.

amplitude: 3, period = π , phase shift = $-\frac{3}{4}\pi$, vertical shift = -3

~~a.~~ $y = \pm 3 \cos \left(\frac{1}{2}\theta - \frac{3}{2}\pi \right) - 3$

c. $y = \pm 3 \cos \left(2\theta - \frac{3}{2}\pi \right) + 3$

~~b.~~ $y = \pm 3 \cos \left(\frac{1}{2}\theta + \frac{3}{2}\pi \right) + 3$

d. $y = \pm 3 \cos \left(2\theta + \frac{3}{2}\pi \right) - 3$

$$\frac{\pi}{1} = \frac{2\pi}{k}$$

$$k\pi = 2\pi$$

$$k = 2$$

C

31. The normal monthly temperatures (°F) for Omaha, Nebraska, are recorded below.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<i>t</i>	1	2	3	4	5	6	7	8	9	10	11	12
Temp.	21°	27°	39°	52°	62°	72°	77°	74°	65°	53°	39°	25°

- a. Write a sinusoidal function that models Omaha's monthly temperature variation.
 b. Use the model to estimate the normal temperature during the month of April.

a. $y = 28 \sin\left(\frac{t}{2}\pi - \frac{\pi}{6}\right) + 49$

b. $y(4) = 49^\circ$

b. $y = 49 \sin\left(\frac{t}{12}\pi + \frac{\pi}{6}\right) + 28$

b. $y(4) = 28^\circ$

c. $y = 28 \sin\left(\frac{t}{6}\pi - \frac{\pi}{2}\right) + 49$

b. $y(4) = 49^\circ$

d. $y = 56 \sin\left(\frac{t}{6}\pi - \frac{\pi}{2}\right) + 28$

b. $y(4) = 56^\circ$

Amp 28

VS 49

P: $\frac{2\pi}{k} = 12 \quad k = \frac{\pi}{6}$

A

32. Write an equation for the given function given the period, phase shift, and vertical shift.

tangent function, period = $\frac{1}{3}\pi$, phase shift = $-\frac{1}{4}\pi$, vertical shift = -5

a. $y = \tan\left(3\theta + \frac{3}{4}\pi\right) - 5$

b. $y = \tan\left(\frac{1}{3}\theta + \frac{3}{4}\pi\right) + 5$

c. $y = \tan\left(\frac{1}{3}\theta - \frac{3}{4}\pi\right) + 5$

d. $y = \tan\left(3\theta - \frac{3}{4}\pi\right) - 5$

$\frac{\pi}{3} = \frac{\pi}{k} \quad \frac{c}{3} = \frac{-\pi}{4}$

$k = 3 \quad 4c = -3\pi$

$c = \frac{-3\pi}{4}$

Write the equation for the inverse of the function.

D

33. $y = \cos 2x$

a. $y = \cos^{-1} 2x$

b. $y = \frac{1}{2} \cos^{-1} 2x$

c. $y = \cos^{-1} \frac{x}{2}$

d. $y = \frac{1}{2} \cos^{-1} x$

$X = \cos ay$

$\cos^{-1} x = 2y$

B

34. $y = \arctan\left(x + \frac{\pi}{2}\right)$

$\tan x = x + \frac{\pi}{2}$

a. $y = \tan x + \frac{\pi}{2}$

b. $y = \tan x - \frac{\pi}{2}$

c. $y = \tan x \left(x - \frac{\pi}{2}\right)$

d. $y = \tan x \left(x + \frac{\pi}{2}\right)$

C 35. $y = \frac{\pi}{4} + \sin x$

$x - \frac{\pi}{4} = \sin y$

a. $y = \frac{\pi}{4} - \text{Arcsin} x$

C c. $y = \text{Arcsin} \left(x - \frac{\pi}{4} \right)$

b. $y = \text{Arcsin} \left(x + \frac{\pi}{4} \right)$

d. $y = \frac{\pi}{4} + \text{Arcsin} x$

C 36. $y = \text{Cos}^{-1}(x - \pi)$

a. $y = \pi - \cos x$

$\cos x = x - \pi$

C c. $y = \pi + \cos x$

b. $y = \cos(x - \pi)$

d. $y = \cos(x + \pi)$

C 37. $y = \text{Arcsin} 3x$

a. $y = \sin \frac{x}{3}$

$\frac{1}{3} \sin x$

C c. $y = \frac{1}{3} \sin x$

b. $y = 3 \sin x$

d. $y = \sin 3x$


C 38. Find the value of $\tan \left(\sin^{-1} \left(\frac{1}{2} \right) \right)$.

a. $\sqrt{3}$

C c. $\frac{\sqrt{3}}{3}$

b. $-\sqrt{3}$

d. $\frac{\sqrt{3}}{3}$



$1^2 + x^2 = 2^2$

$x = \sqrt{3}$

D 39. What basic trigonometric identity would you use to verify that $\tan x \cos x = \sin x$?

a. $\cos x = \frac{1}{\sec x}$

c. $\cos^2 x + \sin^2 x = 1$

b. $\sin x = \frac{1}{\csc x}$

d. $\tan x = \frac{\sin x}{\cos x}$

Chapter 5 starts

B 40. What basic trigonometric identity would you use to verify that $\cot x \sin x = \cos x$?

a. $\cos^2 x + \sin^2 x = 1$

c. $\cos x = \frac{1}{\sec x}$

b. $\cot x = \frac{\cos x}{\sin x}$

d. $\sin x = \frac{1}{\csc x}$

C 41. What basic trigonometric identity would you use to verify that $\frac{\sin^2 x + \cos^2 x}{\cos x} = \sec x$?

a. $\sin x = \frac{1}{\csc x}$

c. $\cos^2 x + \sin^2 x = 1$

b. $1 + \cot^2 x = \csc^2 x$

d. $\cos x = \frac{1}{\sec x}$

C 42. What basic trigonometric identity would you use to verify that $\frac{\sin x + 1}{\sin x} = 1 + \csc x$?

- a. $\sin x = \cos x \tan x$
- b. $\cos^2 x + \sin^2 x = 1$
- c. $\csc x = \frac{1}{\sin x}$
- d. $1 + \cot^2 x = \csc^2 x$

A 43. What basic trigonometric identity would you use to verify that $\sin x \cos x \tan x = 1 - \cos^2 x$?

- a. $\tan x = \frac{\sin x}{\cos x}$
- b. $\cos^2 x + \sin^2 x = 1$
- c. $\sin x = \cos x \tan x$
- d. $1 + \tan^2 x = \sec^2 x$

A 44. Find $\cos x$ if $\sin x \cot x = 4$.

- a. 4
- b. 2

B 45. Find $\cot x$ if $\sin x \cot x \csc x = \sqrt{2}$.

- a. 4
- b. $\sqrt{2}$

C 46. Find $\cos x$ if $\frac{\sin^2 x - 1}{\cos x} = -1$.

- a. -1
- b. 2

D 47. Find $\csc x$ if $\sin x + \cot x \cos x = \sqrt{3}$.

- a. 9
- b. 3
- c. $\frac{\sqrt{3}}{2}$
- d. $\sqrt{3}$

48. Find the exact value of $\cos 15^\circ$.

- a. $\frac{\sqrt{2}}{4}$
- b. $\frac{\sqrt{2} + \sqrt{6}}{4}$
- c. $\frac{\sqrt{2}}{2}$
- d. $\frac{\sqrt{6}}{4}$

49. If α and β are the measures of two first quadrant angles and $\sin \alpha = \frac{4}{5}$ and $\sin \beta = \frac{5}{13}$, find $\sin(\alpha + \beta)$.

- a. $\frac{63}{65}$
- b. $\frac{33}{65}$
- c. $\frac{16}{65}$
- d. $\frac{56}{65}$

50. Which sum or difference identity would you use to verify that $\cos(180^\circ - \theta) = -\cos \theta$?

- a. $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$
- b. $\cos(\alpha - \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$
- c. $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$
- d. $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$

51. Which sum or difference identity would you use to verify that $\sin(90^\circ + \theta) = \cos \theta$?

- a. $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$
- b. $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$
- c. $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$
- d. $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$

#44 hint:
Change $\cot x$ to $\frac{\cos x}{\sin x}$
and simplify
your expression

skip
48-51

(2)